

WARNING

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes eye or skin irritation. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling the concentrate. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, smoking tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep upright with cap tightly closed. Store in original container in areas inaccessible to children.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not refill or reuse this container.

Wrap empty container and put in trash.

Do not reuse empty container.

Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration, if allowed by state and local authorities. Consult federal, state, or local disposal authorities for approved alternate procedure. If burned, stay out of smoke.

BARBICIDE® TB

GERMICIDAL • PSEUDOMONACIDAL • TUBERCULOCIDAL FUNGICIDAL • VIRUCIDAL* • BACTERICIDAL

ACTIVE INGREDIENT:

ortho-Phenylphenol	3.40%
ortho-Benzyl-para-Chlorophenol	3.03%
INERT INGREDIENTS:	93.57%
	100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING

FIRST AID

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** - take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED** - move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **IF SWALLOWED** - call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: In case of an emergency call toll free Chemtrec at 800-424-9300. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**Net Contents: 1 Pint
(16 FL. OZ.) (473 mL)**

Distributed by King Research

7025 W. Marcia Rd • Milwaukee, WI 53223
FOR INDUSTRIAL AND INSTITUTIONAL USE

EPA Reg. No. 34810-31-954
EPA Est. No: 051435-WI-004

GERMICIDAL OR BACTERICIDAL QUALIFICATIONS:

Enterococcus faecalis - Vancomycin Resistant VRE, Methicillin resistant Staphylococcus epidermidis MRSE, Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus-Methicillin resistant MRSA, Staphylococcus aureus **TUBERCULOCIDAL ACTIVITY:** Mycobacterium bovis (BCG) in the presence of 400 ppm hard water (at 20° C for 10 minutes) on hard non-porous surfaces. **FUNGICIDAL ACTIVITY:** AOAC Trichophyton Mentagrophytes **VIRUCIDAL ACTIVITY:** Adenovirus Type 2, Herpes simplex Type 1, Herpes simplex Type 2, Human Immunodeficiency Virus Type 1 HIV-1 (AIDS), Strain: IIIIB, ZeptoMetrix in the presence of 400 ppm hard water and 10% serum in 1 minute. Human Coronavirus, Human Rotavirus, Influenza A2 Virus, Rhinovirus (type 37), Vaccinia virus. BARBICIDE TB may be applied with a cloth, mop or brush, spray or by soaking. For heavily soiled surfaces, thoroughly clean surface prior to disinfecting.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **DO NOT MIX WITH OTHER CLEANERS.** For cleaning and disinfection of hard, non-porous surfaces, use **BARBICIDE TB at a dilution of 1 ounce per gallon of water** for stainless steel, chrome, glass and vinyl. To Disinfect: Wet all surfaces thoroughly. Let stand for 10 minute contact time before wiping or rinsing or allow to air dry. Gross filth and heavy soil must be removed before applying cleaning solution. When used for mopping or from an open pail, solution must be changed at least daily or more often if the solution becomes diluted or soiled. BARBICIDE TB is suitable for cleaning floors in the departments of food processing plants at 1:128 use dilution. Precleaning is required for disinfection of Feline Calicivirus or Norovirus. ***BARBICIDE TB KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HIV-1 (AIDS VIRUS). **PERSONAL PROTECTION:** Wear disposable rubber or latex gloves, gowns, face masks and eye coverings when performing manual cleaning of these surfaces. **CONTACT TIME:** Leave surfaces wet for 1 minute. This contact time will not control other common types of viruses and bacteria. **CLEANING PROCEDURE:** BARBICIDE TB may be used to remove and clean blood and other body fluids from surfaces and objects before disinfection. For lightly soiled areas wipe off visual soil/clean with BARBICIDE TB, and disinfect surface with a second application of BARBICIDE TB. For large spills the contaminated area must be flooded with BARBICIDE TB before cleaning, then decontaminated with fresh BARBICIDE TB. **DISPOSAL OF INFECTIOUS MATERIAL:** Blood and other body fluids must be autoclaved and disposed of according to Federal, State and local regulations for infectious waste disposal. **AREAS OF USE:** Foot spas, Salons (Beauty), Barber Shops. **SALON/BARBER INSTRUMENTS AND TOOLS DISINFECTING BACTERICIDE/VIRUCIDE/FUNGICIDE DIRECTIONS:** Mix 1 oz. (30 mL) of this product per gallon of water (0.25 oz (7.5 mL) of this product per quart of water) to disinfect hard, nonporous instruments and tools. Completely immerse combs, brushes, rollers, manicure and other salon instruments and tools for 10 minutes. For heavily soiled instruments and tools, a preliminary cleaning is required. Rinse thoroughly and dry before use. Fresh solution must be prepared daily or more often if solution becomes diluted or soiled. Note: Plastics may remain immersed until ready to use. Stainless steel instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.



BARBICIDE TB 16 oz. 4/19